



---

Characteristic Numbers for Unoriented Singular  $SG\mathbb{Z}$ -Bordism

Author(s): S. S. Khare and B. L. Sharma

Source: *Proceedings of the American Mathematical Society*, Vol. 61, No. 1 (Nov., 1976), pp. 149-152

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2041682>

Accessed: 04/11/2009 02:35

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

## CHARACTERISTIC NUMBERS FOR UNORIENTED SINGULAR $G$ -BORDISM

S. S. KHARE AND B. L. SHARMA

**ABSTRACT.** We develop the notion of characteristic numbers for unoriented singular  $G$ -manifolds in a  $G$ -space,  $G$  being a finite group, and prove their invariance with respect to unoriented singular  $G$ -bordism.

Thom [5] gave the notion of Stiefel Whitney numbers and Pontrjagin numbers of a manifold  $M^n$  and proved its invariance with respect to bordism. Chung N. Lee and Arthur Wasserman [4] developed these notions for  $G$ -manifolds. In this note we have developed these notions for unoriented singular principal  $G$ -manifolds in a  $G$ -space,  $G$  being a finite group, and proved their invariance with regard to unoriented singular  $G$ -bordism.

**1. Characteristic numbers.** Let  $X$  be a finite CW-complex with free action of  $G$ ,  $G$  being a finite group, and  $X/G$  be again a finite CW-complex. Let  $h^*$  be an equivariant cohomology theory and  $h_*$  be the associated equivariant homology theory [1]. Let  $h^* = H^* \circ A$  and  $h_* = H_* \circ A$ , where  $A$  is a functor from the category of  $G$ -spaces and equivariant maps to the category of topological spaces and continuous maps,  $H^*$  is the singular cohomology theory and  $H_*$  is the associated singular homology theory. Let

$$\langle , \rangle : h^*(X; G) \otimes_{H^*(\text{pt.})} h_*(X; G) \rightarrow H_*(\text{pt.})$$

be the Kronecker pairing.

Let us assign to each compact  $G$ -manifold  $W$ , a class

$$[W, \partial W] \in h_*(W, \partial W; G)$$

such that

- (a)  $[W_1 \cup W_2, \partial W_1 \cup \partial W_2] = [W_1, \partial W_1] + [W_2, \partial W_2]$ ,
- (b)  $\partial_*[W, \partial W] = [\partial W]$ .

Suppose  $[M^n, f; G]$  is an element of unoriented bordism group  $\mathfrak{N}_n(\check{X}; G)$  [3] and  $x \in h^*(B(O, G)_n; G)$ ,  $B(O, G)_n$  being the classifying space for  $G$ -vector bundles of dimension  $n$ . Then the  $x$ -characteristic number of the map  $f: M^n \rightarrow X$  associated with an element  $a^m \in h^m(X; G)$  is defined to be

Received by the editors August 22, 1974.

AMS (MOS) subject classifications (1970). Primary 57D85.

Key words and phrases.  $G$ -bordism, equivariant homology, equivariant cohomology.

Copyright © 1977, American Mathematical Society

$\langle \tau_{M^n}^*(x) f^*(a^m), [M] \rangle$ , where  $\tau_{M^n}: M^n \rightarrow B(O, G)_n$  is the tangent map.

In particular, let the equivariant cohomology  $h^*$  be given by  $h^*(X; G) = H^*((E_G \times X)/G; \mathbf{Z}_2)$  and  $h_*$  be the associated equivariant homology, i.e.  $h_*(X; G) = H_*((E_G \times X)/G; \mathbf{Z}_2)$ , where the action of  $G$  on  $E_G \times X$  is given by  $g(e, x) = (ge, gx)$ ,  $E_G$  being the total space of the universal  $G$ -bundle. Consider the map  $q: X/G \rightarrow (E_G \times X)/G$  given by  $q([x]) = [\bar{\alpha}(x), x]$ , where  $\bar{\alpha}$  is the map given by the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{\bar{\alpha}} & E_G \\ \downarrow & & \downarrow \\ X/G & \xrightarrow{\alpha} & BG \end{array}$$

The map  $q$  is homotopy equivalence. Thus

$$h^*(X; G) \overset{q^*}{\approx} H^*(X/G; \mathbf{Z}_2) \quad \text{and} \quad h_*(X; G) \overset{q_*^{-1}}{\approx} H_*(X/G; \mathbf{Z}_2).$$

Therefore  $h_*(M^n; G) \approx H_*(M^n/G; \mathbf{Z}_2)$  has a topological class, say  $\sigma_n$ , in dimension  $n$ .

**2. Invariance of characteristic numbers.** Throughout the section we will be considering equivariant cohomology  $h^*$  to be

$$h^*(X; G) = H^*((E_G \times X)/G; \mathbf{Z}_2)$$

and equivariant homology  $h_*$  to be

$$h_*(X; G) = H_*((E_G \times X)/G; \mathbf{Z}_2).$$

**THEOREM 2.1.** *If  $[M^n, f; G] \in \mathfrak{N}_n(X; G)$  is zero then all the  $x$ -characteristic numbers of the map  $f: M^n \rightarrow X$  associated with every  $a^m \in h^m(X; G)$  are zero.*

**PROOF.** Since  $[M^n, f; G] \in \mathfrak{N}_n(X; G)$  is zero,  $\exists$  an  $(n + 1)$ -dimensional compact principal  $G$ -manifold  $W^{n+1}$  and an equivariant map  $F: W^{n+1} \rightarrow X$  with  $\partial W^{n+1} = M^n$  and  $F/M^n = f$ . Let  $\omega_{n+1} \in h_{n+1}(W^{n+1}, \partial W^{n+1}; G)$  be the topological class of  $W^{n+1}$ . Then  $\partial_* \omega_{n+1} = \sigma_n$ . We have the following commutative diagram:

$$\begin{array}{ccc} h^*(B(O, G)_n; G) & \xrightarrow{\tau_{M^n}^*} & h^*(M^n; G) \\ \uparrow j^* & & \uparrow i^* \\ h^*(B(O, G)_{n+1}; G) & \xrightarrow{\tau_{W^{n+1}}^*} & h^*(W^{n+1}; G), \end{array}$$

where  $j: B(O, G)_n \rightarrow B(O, G)_{n+1}$  is the map classifying  $\mu_n \oplus 1$ ,  $\mu_n \rightarrow B(O, G)_n$  being the universal  $G$ -vector bundle. Also we have

$$\begin{aligned} h^*(B(O, G)_n; G) &= H^*((E_G \times B(O, G)_n)/G; \mathbf{Z}_2) \\ &= H^*(BG \times BO_n; \mathbf{Z}_2) \quad [6] \\ &= H^*(BG; \mathbf{Z}_2) \otimes H^*(BO_n; \mathbf{Z}_2) \end{aligned}$$

and

$$h_*(B(O, G)_n; G) = H_*(BG; \mathbf{Z}_2) \otimes H_*(BO_n; \mathbf{Z}_2).$$

Thus the map  $j^*$  is a surjection. Therefore for every  $x \in h^*(B(O, G)_n; G)$ ,  $\exists y \in h^*(B(O, G)_{n+1}; G)$  such that  $j^*(y) = x$ . Therefore

$$\begin{aligned} \langle \tau_{M^n}^*(x) f^*(a^m), \sigma_n \rangle &= \langle \tau_{M^n}^* j^*(y) f^*(a^m), \sigma_n \rangle = \langle i^* \tau_{W^{n+1}}^*(y) i^* F^*(a^m), \sigma_n \rangle \\ &= \langle \tau_{W^{n+1}}^*(y) F^*(a^m), i_* \partial_*(\omega_{n+1}) \rangle = 0. \end{aligned}$$

This completes the proof of the theorem.

Consider now the map  $\mu : \mathfrak{N}_*(X; G) \rightarrow h_*(X; G)$  defined as  $\mu([M^n, f; G]) = q_* \bar{f}_*(\bar{\sigma}_n)$ , where  $\bar{f}$  is the map given by the following commutative diagram:

$$\begin{array}{ccc} M^n & \xrightarrow{f} & X \\ \downarrow & & \downarrow \\ M^n/G & \xrightarrow{\bar{f}} & X/G \end{array}$$

$\bar{\sigma}_n \in H_n(M^n/G; \mathbf{Z}_2)$  being the fundamental class and let  $\bar{\mu} : \mathfrak{N}_*(X/G) \rightarrow H_*(X/G; \mathbf{Z}_2)$  be the map defined by  $\bar{\mu}([N^n, g]) = g_*(\bar{\sigma}'_n)$ , where  $\bar{\sigma}'_n \in H_n(N^n; \mathbf{Z}_2)$  is the fundamental class. Suppose  $\phi_* : \mathfrak{N}_*(X; G) \rightarrow \mathfrak{N}_*(X/G)$  is the isomorphism [3] defined as  $\phi_*([M^n, f; G]) = [M^n/G, \bar{f}]$ . Then  $\mu = q_* \bar{\mu} \phi_*$  and, therefore,  $\mu$  is an epimorphism, since  $\bar{\mu}$  is so [2]. For every  $a \in h_*(X; G)$ , we select  $[M^n, f; G] \in \mathfrak{N}_*(X; G)$  such that  $\mu([M^n, f; G]) = a$ . We define the  $\mathfrak{N}$ -module structure on  $h_*(X; G)$  by

$$[V^m]a = \mu[M^n \times V^m, f'; G],$$

for every  $[V^m] \in \mathfrak{N}$ , where the action of  $G$  on  $M^n \times V^m$  is defined as  $g(x, y) = (gx, y)$  and  $f' : M^n \times V^m \rightarrow X$  is defined as  $f'(x, y) = f(x)$ . Thus  $h_*(X; G) \otimes \mathfrak{N}$  is a  $\mathfrak{N}$ -module. Let  $\{C_{n,i}\}$  be the additive base of  $h_*(X; G)$ . Let  $[M_i^n, f_i; G] \in \mathfrak{N}_*(X; G)$  with  $\mu([M_i^n, f_i; G]) = C_{n,i}$ . We define  $h : h_*(X; G) \otimes \mathfrak{N} \rightarrow \mathfrak{N}_*(X; G)$  by  $h(C_{n,i} \otimes 1) = [M_i^n, f_i; G]$ .

**THEOREM 2.2.** *The map  $h : h_*(X; G) \otimes \mathfrak{N} \rightarrow \mathfrak{N}_*(X; G)$ , defined as above is an isomorphism.*

**PROOF.** We have the following commutative diagram:

$$\begin{array}{ccc} h_*(X; G) \otimes \mathfrak{N} & \xrightarrow{h} & \mathfrak{N}_*(X; G) \\ \downarrow q_*^{-1} \otimes 1_{\mathfrak{N}} & & \downarrow \phi_* \\ H_*(X/G; \mathbf{Z}_2) \otimes \mathfrak{N} & \xrightarrow{\bar{h}} & \mathfrak{N}_*(X/G) \end{array}$$

where  $\bar{h} : H_*(X/G; \mathbf{Z}_2) \otimes \mathfrak{N} \rightarrow \mathfrak{N}_*(X/G)$  is defined as  $\bar{h}(\bar{C}_{n,i} \otimes 1) = [M_i^n/G; \bar{f}_i]$ , where  $\bar{C}_{n,i} = q_*^{-1}(C_{n,i})$ . We already know that  $\bar{h}$  is an isomorphism [2] and, therefore, so is  $h$ .

The above theorem gives the converse of Theorem 2.1 given as below.

**THEOREM 2.3.** *If all the characteristic numbers of an element  $[M^n, f; G] \in \mathfrak{R}_*(X; G)$  are zero, then  $[M^n, f; G] = 0$ .*

**PROOF.** Let  $\mu([M^n, f; G]) = C_n \in h_*(X; G)$  and  $q_*^{-1}(C_n) = \bar{C}_n \in H_*(X/G; \mathbf{Z}_2)$ . Therefore  $\bar{f}_*(\bar{\sigma}_n) = \bar{C}_n$ . Suppose  $\{C_{n,i}\}_{i \in I}$  is an additive base of  $h_n(X; G)$  and  $C^{n,j} \in h^n(X; G)$  is the cohomology class dual to  $C_{n,i}$  in the sense  $\langle \bar{C}^{n,j}, \bar{C}_{n,i} \rangle = \delta_{ij}$ , where  $q_*^{-1}(C_{n,i}) = \bar{C}_{n,i}$  and  $q^*(C^{n,j}) = \bar{C}^{n,j}$ . Let  $C_n = \sum_{i \in S} \pm C_{n,i}$ ,  $S$  being a finite subset of  $I$ . Then if  $C^n = \sum_{i \in S} \pm C^{n,i}$ , by hypothesis the  $x$ -characteristic number of  $[M^n, f; G]$  associated with  $C^n \in h^n(X; G)$  is zero, that means taking  $x$  to be unit class of  $h^*(B(O, G)_n; G)$ ,

$$\langle f^*(C^n), [M] \rangle = 0, \quad \text{or} \quad \langle f^*(C^n), q_*[\bar{\sigma}_n] \rangle = 0,$$

or

$$\langle (q^*)^{-1}(\bar{C}^n), \bar{f}_* q_*[\bar{\sigma}_n] \rangle = 0, \quad \text{or} \quad \langle (q^*)^{-1}(\bar{C}^n), q_* \bar{f}_*[\bar{\sigma}_n] \rangle = 0,$$

by the following commutative diagram

$$\begin{array}{ccc} h_n(M^n; G) & \xrightarrow{f_*} & h_n(X; G) \\ \downarrow q_*^{-1} & & \downarrow q_*^{-1} \\ H_n(M^n/G; \mathbf{Z}_2) & \xrightarrow{\bar{f}_*} & H_n(X/G; \mathbf{Z}_2) \end{array}$$

Therefore  $\langle (q^*)^{-1}(\bar{C}^n), q_*(\bar{C}_n) \rangle = 0$ , which implies that  $\langle \bar{C}^n, \bar{C}_n \rangle = 0$ , showing that  $\bar{C}_n = 0$ . Also it is easy to see that  $h(C_n \otimes 1) = [M^n, f; G]$ . Since  $h$  is an isomorphism and  $C_n = 0$ ,  $[M^n, f; G] = 0$ , which completes the proof of the theorem.

Theorems 2.1 and 2.3 give the invariance of characteristic numbers with regard to unoriented singular principal  $G$ -bordism.

REFERENCES

1. G. E. Bredon, *Equivariant cohomology theories*, Lecture Notes in Math., vol. 34, Springer-Verlag, Berlin and New York, 1967. MR 35 #4914.
2. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Academic Press, New York; Springer-Verlag, Berlin, 1964. MR 31 #750.
3. S. S. Khare and B. L. Sharma, *Equivariant bordism, cobordism and duality theorem* (communicated).
4. C. N. Lee and A. G. Wasserman, *Equivalent characteristic numbers* (Proc. Second Conf. Compact Transformation Groups, 1971), Part I, Lecture Notes in Math., vol. 298, Springer-Verlag, Berlin, 1972, pp. 191–216. MR 51 #1853.
5. R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment Math. Helv. 28 (1954), 17–86. MR 15, 890.
6. T. tom Dieck, *Faserbündel mit Gruppenoperation*, Arch. Math. (Basel) 20 (1969), 136–143. MR 39 #6340.

DEPARTMENT OF MATHEMATICS, NORTH-EASTERN HILL UNIVERSITY, RITA VILLA, LOWER LACHAUMIERE, SHILLONG-793 001, MEGHALAYA, INDIA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALLAHABAD, ALLAHABAD (U.P.), INDIA