

Seasonal Fluctuation of Different Edaphic Microarthropod Population Densities in Relation to Soil Moisture and Temperature in a Pine, *Pinus* *kesiya* Royle Plantation Ecosystem*

by

M. Vikram Reddy**

ABSTRACT. – Seasonal fluctuations of soil and litter microarthropod populations in a pine, *Pinus kesiya* Royle plantation of North Eastern India were investigated between November 1976 and November 1977. Three major groups were recognized: (a) Collembola, (b) Acarina and (c) miscellaneous. Collembola was the most abundant group and was dominated by *Isotoma trispinata* (MacGillivray). The total microarthropod density ranged from 26,800 per m² to 145,200 per m². Collembola densities ranged from 10,000 to 121,200 per m², Acarina densities ranged from 8,800 to 41,600 per m², and the miscellaneous group ranged from 1,200 to 6,400 per m². Soil moisture was positively correlated with total arthropod, Collembola and Acarina densities. Soil temperature was positively correlated only with Acarina. Densities of Collembola and Acarina were negatively correlated.

INTRODUCTION

Edaphic arthropods living in forest/plantation soils play a significant role in litter decomposition (Crossley, 1978). Their densities and their activity, however, vary with season. Studies to determine seasonal patterns of arthropod populations in tropical and sub-tropical forests have been few, particularly in the Indian subcontinent (Singh and Singh, 1975; Prabhoo, 1976). Little is known about the relationships between arthropod populations and the abiotic environment of Indian pine forests and plantations. The objective of this study was to determine how the seasonal abundance of major groups of soil and litter microarthropod populations might be related to soil moisture and temperature in a pine (*Pinus kesiya* Royle) plantation near Shillong (Meghalaya), North Eastern India.

* The work described in this paper was carried out while the author was at the Department of Zoology, North Eastern Hill University, Shillong, Meghalaya (India).

It was presented at the Ninth International Biometeorological Congress, 23rd September-1st October, 1981, Osnabrück and Stuttgart-Hohenheim, FRG.

** Current address is: Agricultural College, NEHU, Medziphema-797106, Nagaland, INDIA.

MATERIALS AND METHODS

The pine plantation studied in the present report has been described by Reddy (1981a). Monthly soil and litter samples were taken with a 5 x 5 x 10 cm soil sampler during the early morning hours from November 1976 to November 1977. On each occasion ten random samples were taken and carefully placed into the polythene bags for transport to the laboratory. Within five to six hours of collection, the samples were placed into a series of modified Tullgren funnels and processed for six to seven days. Microarthropods were collected in 80% ethyl alcohol. In addition to soil and litter samples, soil temperature was monitored at a 5 cm depth and an extra set of soil samples was collected for a gravimetric determination of soil moisture (Piper, 1944). Finally, correlation coefficients were determined between monthly mean densities of the various microarthropod groups and monthly soil temperatures and moisture readings.

RESULTS

Microarthropods from samples were grouped into the following categories: (a) Collembola, (b) Acarina and (c) miscellaneous. The latter group included Isopoda, Thysanoptera, Hymenoptera, Araneidae, Protura Diplura, Chilopoda, Diplopoda, Symphyla, Pauropoda, microcoleopteran adults and their larvae and Dipteran larvae. The Collembola being the most dominant group among the microarthropods at this site, was represented by five families (Isotomidae, Entomobryidae, Onychiuridae, Sminthuridae and Hypogastridae) and dominated by *Isotoma trispinata* (MacGillivray). Acarina was represented by Prostigmata, Mesostigmata and Cryptostigmata being dominated by Prostigmata. Astigmata was absent. The total arthropod population consisted of 63.16% Collembola, 31.79% Acarina and 5.05% miscellaneous species.

Seasonal fluctuations of total microarthropod population ranged from 26,800 per m² in September of 1977 to 145,200 per m² in July of 1977 (Fig. 1). The density per m² of Collembola ranged from 10,000 in January of 1977 to 121,200 in July of 1977. Acarina ranged from 8,800 in May of 1977 to 41,600 per m² in November of 1977. Finally, the miscellaneous group ranged from 1200 in May of 1977 to 6,400 per m² in November of 1976. Collembola dominated the total population from April to October, 1977 whereas Acarina dominated during November 1976 to March 1977 and again in November of 1977. The correlation coefficient between the monthly abundance of Collembola and Acarina was negative ($r = -0.3$, $P < 0.2$).

Mean soil temperature taken at the various sample times ranged from 16°C in December of 1976 to 24°C in August of 1977. Soil moisture ranged from 12% in January to 36% in July of 1977 (Fig. 1).

The correlation coefficient for microarthropods and soil moisture were positive and significant for total arthropods, Collembola and Acarina (Table 1). The correlation coefficient for microarthropods and soil temperature were significant and positive only for Acarina.

DISCUSSION

The Collembola group, which was dominated by *Isotoma trispinata*, and the Acarina group comprised 95% of the total microarthropods in this study. Similar findings have




Fig. 1. Histogram showing seasonal population abundance of different soil mesofaunal groups of a pine wood (*Pinus kesiya-royle*) Soil (N.E. India) in relation to soil moisture and temperature.

Table 1. Correlation coefficients among various groups of edaphic microarthropods, and soil moisture and temperature in a pine, *Pinus kesiya* plantation ecosystem. (n = 13).

Variable "X"	Variable "Y"	Coefficient correlation (r)	Computed value of "t"	Level of significance
Total arthropods	Moisture	+ 0.5693	2.296	P<0.05
Collembola	Moisture	+ 0.7251	3.492	P<0.01
Acarina	Moisture	+ 0.6274	2.653	P<0.05
Miscellaneous group	Moisture	+ 0.1500	0.510	P<0.7 ^a
Total arthropods	Temperature	+ 0.1295	0.433	P<0.7 ^a
Collembola	Temperature	+ 0.3262	1.144	P<0.3 ^a
Acarina	Temperature	+ 0.6178	2.606	P<0.05
Miscellaneous group	Temperature	- 0.2400	0.830	P<0.5 ^a

a = Not significant

been reported for summer populations in tea nursery beds (Reddy, 1981b) and rose-garden soil (Mukharjii and Singh, 1970). However, Fujikawa (1970) and Price (1973) reported that Acarina was a dominant group in their studies of forest soil. Variation in population densities have also been reported (Poole, 1961; Price, 1973). Crossley and Bohnsack (1960) reported a microarthropod density of 102,00 per m² in a pine forest in Tennessee. Such variations both in species composition and density are probably due to differences in the physical and biological conditions of the various study sites (Im-

madate and Kira, 1964). That the physical environment is important in this regard is implied by the positive correlations between Collembola and Acarina and soil moisture and Acarina and temperature. Positive correlations between microarthropods and soil moisture have also been reported by others (Price, 1973; Nijima, 1975; Kaczmarek, 1975; Choudhuri, Hazra and Roy, 1978). With the exception of Acarina, the correlations between microarthropod densities and temperature were not significant. Durant and Richard (1966) observed no correlations between density and temperature in their study.

The fact that the correlation between Collembola and Acarina were negative may indicate an antagonistic relationship between them, possibly due to predation of mites on Collembola (Nijima, 1971; Reddy, 1980).

I. trispinata which was most dominant and recorded through out the year may be designated as an indicator species of the system.

ACKNOWLEDGEMENTS

The author is thankful to Prof. Dr. R. G. Michael and Prof. Dr. P. S. Ramakrishnan, School of Life-Sciences, NEHU for providing necessary facilities, and to Dr. J. R. B. Alfred for encouragement. This research was funded by a grant from University Grants Commission, New Delhi.

REFERENCES

- CHOUDHURI, D. K., HAZRA, A. K. and ROY, S. (1978): Soil factors governing the distribution of Collembola (Insecta) in the grave-yard of Berhampore, Murshidabad district (West Bengal). In: Soil Biology and Ecology in India: C. A. Edwards, and G. K. Veeresh (eds.), UAS (Bangalore) Techn. Series No. 22: 161-172.
- CROSSLEY, D. A., Jr. (1978): The role of terrestrial saprophagous arthropods in forest soils: Current status of concepts. In: The Role of Arthropods in Forest Ecosystems: W. J. Mattson (ed.), Springer-Verlag, New York, 49-56.
- CROSSLEY, D. A., Jr. and BOHNSACK, K. K. (1960): Long term ecological study in the Oak Ridge area: 3. The Oribatid mites fauna in pine litter. Ecology, 41: 628-638.
- DURANT, J. A. and RICHARD, F. C. (1966): Some arthropods of forest floor in pine and hardwood forests in the South Carolina Piedmont region. Ann. Entomol. Soc. Amer., 59: 202-207.
- FUJIKAWA, T. (1970): Distribution of soil animals in three forests of Northern Hokkaido. II Horizontal and vertical distributions of Oribatid mites (Acarina: Cryptostigmata). Appl. Ent. Zool., 5: 208-212.
- IMMADATE, G. and KIRA, T. (1964): Notes on the soil microarthropods collection made by the Thai-Japanese Biological Expedition 1961-62. Description of collecting sites, group contents and ecological characters of 180 samples from South East Asia. Fauna and Flora Res. Soc., Kyoto: 81-111.
- KACZMAREK, M. (1975): An analysis of Collembola communities in different pine forest environments. Ecol. Pol., 23: 265-293.
- MUKHARJI, S. P. and SINGH, J. (1970): Seasonal variation in the densities of a soil

- arthropod population in a rose-garden at Varanasi (India). *Pedobiologia*, 10: 442-446.
- NIJIMA, K. (1971): Seasonal changes in Collembolan populations in a warm temperate forest of Japan. *Pedobiologia*, 11: 11-26.
- NIJIMA, K. (1975): Seasonal changes in Collembolan populations in a warm temperate forest of Japan. II. Population dynamics of the dominant species. *Pedobiologia*, 15: 40-52.
- PIPER, C. S. (1944): *Soil and Plant Analysis*. The University of Adelaide, Adelaide, 1-368 pp.
- POOLE, T. B. (1961): An ecological study of the Collembola in a coniferous forest soil. *Pedobiologia*, 1: 113-137.
- PABHOO, N. R. (1976): Soil microarthropods of a virgin forest and adjoining tea fields in the Western Ghats in Kerela – A brief ecological study. *Oriental insects*, 10: 435-442.
- PRICE, D. W. (1973): Abundance and vertical distribution of microarthropods in the surface layers of California pine forest soil. *Hilgardia*, 42: 121-148.
- REDDY, M. V. (1980): Arthropod faunal structure of soil surface understone habitats at Berhampur, Orissa. *Newsletter Soil Biology & Ecology*, 1: 18-19.
- REDDY, M. V. (1981a): Microarthropods and the rate of needle litter disappearance in a pine plantation ecosystem of North Eastern India. *Pedobiologia*, 22: 339-344.
- REDDY, M. V. (1981b): Faunal composition of summer population of soil microarthropods of tea nursery beds at Nayabunglow, North Eastern India. *Indian J. Ecol.*, 8: 108-111.
- SINGH, J. and SINGH, U. R. (1975): An ecological study of soil microarthropods upon soil and litter of a tropical deciduous forest of Varanasi (India). *Trop. Ecol.*, 16: 81-85.