

Effect of Acidic Coal Mine Drainage on Soil Fertility Status of Rice Soils of Jaintia Hills District of Meghalaya

B. Goswami¹, O. P. Singh²; K. K. Satapathy¹ and U. S. Saikia^{1*}

¹Division Of Agricultural Engineering, ICAR Research Complex For NEH Region, Umiam, 793 103, Meghalaya

²Centre For Environmental Studies, North Eastern Hill University, Shillong, Meghalaya

*Corresponding author

Received in January, 2008

Revised and accepted in May, 2008

Analysis of soil samples collected from rice fields of three different coal mining areas of Jaintia Hills district, Meghalaya, namely, Dkhiah, Lad Rymbal and Bapung revealed that soils were highly acidic (pH 3.3-4.1) with inadequate availability of N (210-272 kg/ha), P₂O₅ (21-33 kg/ha), K₂O (145-210 kg/ha) and organic carbon (0.21-0.67 %). The soil fertility status was found to be not congenial to sustain plant growth and development in long term.

Key words : Mine drainage, Nutrient, pH

INTRODUCTION

The Jaintia Hills district of Meghalaya is the most extensively exploited in terms of coal, among all the districts of the state¹. Coal mining operations in Jaintia Hills are small scale ventures controlled by individuals who own the land. Coal extraction is done following primitive mining method known as 'rat hole' mining. This extensive surface excavation of sloping land damages the ecosystem within the periphery of mined area².

Agriculture is the main occupation of the people of Jaintia Hills. Rice is the main crop of this region occupying 50.5 % of total net sown area³. But during recent years large number of paddy fields are left barren by the farmers due to degradation of fields through deposition of coal particles⁴ and contamination of soils due to inundation by highly acidic mine water⁵. Rectification of such problem is possible only if inherent properties of such soils with respect to crop growth is studied. This paper attempts to focus on some basic properties of soils of paddy fields located in different coal mining sites of Jaintia Hills District.

Experimental

STUDY AREA AND METHODS

Jaintia Hills district is situated in the eastern part of the state of Meghalaya and lies between 25°5' to 25°4'

N latitudes and between 91°51' to 92°45'E longitudes. Coal fields of Jaintia Hills are small and spread out in different patches, namely, Bapung, Lakadong, Lumshong, Malwar-Muslang Lamare, Mutang, Sutnga, Jarain-Tkentalang and Ioksi. Although there is large-scale exploitation of coal in the district but for the purpose of study three representative coal mining sites, namely, Lad Rymbal, Bapung and Dkhiah have been selected.

Surface soil samples (0-15 cm depth) collected from paddy fields were mixed, sieved and homogenized for determining various physico-chemical characteristics. Geographical locations of the sampling sites are depicted in Table 1.

Soil pH was determined by glass electrode pH meter (1:2.5 :: Soil: water). Organic carbon⁶, available N⁷, available P and available K⁸ were determined by following standard chemical analytical procedures.

RESULTS AND DISCUSSION

Soil pH and organic carbon status of experimental soils have been presented in Table 2. Soil pH of coal mining sites ranged from 3.3 to 4.1 indicating strongly acidic nature. This is below the optimum pH (5.5 to 6.0) required for rice⁹. Among all the sites pH was lowest in

Table1: Geographical locations of sampling sites

Location	Altitude (above msl*)	Latitude (N)	Longitude (E)
Lad Rymbal	1,200 m	25°22'14"	92°19'4"
Dkhlah	1,200 m	25°22'18"	92°21'32"
Bapung	1,262 m	25°23'45"	92°18'24"

*msl: mean sea level

Bapung. The upper range of organic carbon for proper growth and development of crop is 0.75%¹⁰. The data shows that this limit is not attained at all sampling sites. Moreover both pH and organic carbon of experimental soils were lower than pH (4.7-5.4) and organic carbon (0.9-7.41%) of rice soils of non-mining areas of Jaintia Hills¹¹. Low level of organic carbon (0.21 to 0.67%)

21-33 and 145-210 kg/ha, respectively. Availability of all these nutrients were in medium range which is not sufficient to sustain optimum growth and development of crops. Moreover availability of N, P₂O₅ and K₂O of Bapung soil were lowest among all the sampling sites as compared to rice soils of unmined areas of Jaintia Hills (246-268 kg/ha N, 21-32 kg/ha P, 164-169 kg/ha K)¹¹.

Table 2: Soil pH and organic carbon of experimental soils

Soil Parameters	Dkhlah		Lad Rymbal		Bapung	
	Mean*	SD	Mean	SD	Mean	SD
pH	3.80	0.36	4.10	0.38	3.30	0.26
Organic C (%)	0.53	0.08	0.67	0.15	0.21	0.05

*Mean of three samples

indicates poor soil physical properties which ultimately results in poor growth and development of crop grown in such soils¹².

Table 3 indicates that available N, P₂O₅ and K₂O in all the experimental soils were in the ranges of 210-272,

Lower availability of nutrients to plants is responsible for decreased vegetative growth which causes reduced development of photosynthetic area and thereby lower translocation of photosynthates towards sinks which ultimately results in poor growth and yield of crops¹³.

Table 3: Available major nutrients of experimental soils

Soil Parameters	Dkhlah		Lad Rymbal		Bapung	
	Mean*	SD	Mean	SD	Mean	SD
Available N (kg/ha)	247.43	32.86	272.87	23.38	210.73	19.81
Available P ₂ O ₅ (kg/ha)	33.23	4.61	25.40	5.60	21.53	1.62
Available K ₂ O (kg/ha)	176.50	20.79	210.13	11.87	145.00	11.36

*Mean of three samples

DISCUSSION

The results of the investigation suggest that the soils collected from paddy fields located in coal mining sites of Jaintia Hills district, Meghalaya were highly acidic with inadequate availability of major plant nutrients and organic carbon. The soils can not sufficiently support proper growth and development of crops. It needs immediate attention before the soil productivity is deteriorated irreversibly.

REFERENCE

1. Anonymous, In: *Technical Report of the Directorate of Mineral Resources*, Government of Meghalaya, 2003.
2. Das Gupta S, Tiwari B K and Tripathy R S, Coal mining in Jaintia Hills, Meghalaya: An Ecological Perspective, In: *Jaintia Hills, A Meghalaya Tribe: Its Environment, Land and People*. (Eds. P M Passah and A S Sarma), Reliance Publishing House, New Delhi, 2002, 121.
3. Anonymous, *Meghalaya socio-economic review*, Directorate of Economics and Statistics, Government of Meghalaya, Shillong, India, 2003.
4. Nath M and Ahmed M, *Mining Environment: Problems and Remedies* (Eds. Singh O P), Regency Publications, New Delhi, India, 2005, 70.
5. Swer S and Singh O P, *ENVIS Bulletin* 11(2), 2003, 26.
6. Walkley A and Black I A, *Soil Science* 34, 1934, 29.
7. Subbiah B V and Asija G L, *Current Sc*, 25, 1956, 259.
8. Bray R H and Kurtz L T, *Soil Sc*, 59, 1954, 39.
9. Barbosa Filho M P and Yamada, *Bollor Cropn International*, 10, 2002, 43.
10. Jackson M L, *Soil chemical analysis*. Prentice Hall of India Pvt. Ltd., New Delhi, India, 1973.
11. Venkatesh M S, Kailash Kumar, Rauchaudhuri M, Datta M and Sharma U C, Soil fertility and crop responses to nutrients, In: *Soils and their management in North east India* (Eds. U C Sharma, M Datta and J S Samra), ICAR Research Complex for NEH Region, Umiam, Meghalaya Publication, 2006, 280.
12. Channabasavanna A S, Itnal C J and Patol S G, *Indian J Agron*, 47, 2001, 1.
13. Parihar S S, *Ann Agril Res*, 25 (3), 2004, 408.