

STUDIES ON THE VESICULAR ARBUSCULAR MYCORRHIZA OF
PADDY IN MEGHALAYA

Ajay Kumar Singh

THESIS
SUBMITTED
IN
FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

To



DEPARTMENT OF BOTANY
SCHOOL OF LIFE SCIENCES
NORTH-EASTERN HILL UNIVERSITY

SHILLONG - 793014

INDIA

DECEMBER, 1989



COMPUTER

Grams : NEHU

Phone : Off 23390

North - Eastern



Hill University

PROFESSOR R. R. MISHRA
M. Sc., Ph. D., F. N. A. Sc., F. N. I. E.

DEPARTMENT OF BOTANY
SCHOOL OF LIFE SCIENCES
SHILLONG - 793014

I certify that the thesis entitled "Studies on the Vesicular- arbuscular mycorrhiza of paddy in Meghalaya", submitted by Ajay Kumar Singh, for the degree of Doctor of Philosophy, embodies the record of original investigation carried out by him under my supervision. He has been duly registered and the thesis presented is worthy of being considered for the award of the Ph.D. degree. This work has not been submitted for any degree of any other University.

Date :

Place : Shillong

(R.R.MISHRA)

Signature of the Supervisor

Forwarded
Y.S. Chakraborty
21.2.90
Head
Department of Botany
School of Life Sciences
N.E.H.U., Shillong-793014

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	6
MATERIALS AND METHODS	29
RESULTS	
1. Status of vesicular-arbuscular mycorrhiza in paddy	51
2. Effect of vesicular-arbuscular mycorrhiza on growth and P uptake of phosphorus deficiency tolerant and susceptible paddy varieties (RCPL 101 and RCPL 104 respectively) under different soil phosphorus level	55
3. Effect of vesicular-arbuscular mycorrhiza on growth and nutrient uptake in paddy at different soil nitrogen and phosphorus levels	62
4. Effect of vesicular-arbuscular mycorrhiza on growth of paddy under different water stress conditions	71
5. Interaction of vesicular-arbuscular mycorrhizal fungi with rhizospheric fungi of paddy	83
DISCUSSION	87
SUMMARY	111
LITERATURE CITED	118

ACKNOWLEDGEMENTS

It is my great pleasure to have this opportunity of expressing my heartfelt sense of gratitude to Professor R.R. Mishra, Ph.D., F.N.A.Sc., F.N.I.E., Head of the Department of Botany, School of Life Sciences, North-Eastern Hill University, Shilong, for his able guidance and constant encouragement throughout the period of the present research work.

This work was made possible because of valuable and mature counselling of Dr. G.D. Sharma and Dr. B.K. Tiwari, both Reader in the Department of Botany, NEHU, Shillong, during research work and writing of the manuscript. I am grateful to Dr. N.D. Mazumdar, Plant Breeder, for providing paddy seeds, Mr. Munna Ram, Soil Scientist, ICAR Research Complex for N.E.H. Region, Shillong, for extending necessary facilities and help during some of the chemical analysis, Dr. D.S. Hayman, Rothamstead Experimental Station, Harpenden, Hertfordshire, U.K., for providing cultures of vesicular-arbuscular fungi - Glomus mosseae and G. etunicatum and Dr. K.D. Sharma, ICAR Research Complex for N.E.H. Region, Shillong for the culture of G. caledonium.

I sincerely thank Mr. D.K. Jha, Mr. S. Jyrwa, Mr. C.S. Rao, Mr. S.K. Sharma, Dr. A.N. Raghuwansi and others in the laboratory for helping in various ways.

At last, but not least, I shall like to extend my sincere thanks to Miss R. Sunabi and Mr. B.K Das, who deserve special mention for typing and photography respectively.

I am greatly indebted to my family members for constant inspiration during the study period.

The Director, ICAR Research Complex for N.E.H. Region, Shillong is also duly acknowledged for granting permission to take up the study.

Shillong,
October , 1989.

Ajay Kumar Singh
(Ajay Kumar Singh)

INTRODUCTION

Paddy is the most important staple crop of Meghalaya. Due to high rainfall during kharif season, in addition to transplanted lowland paddy cultivation, the direct sown upland paddy cultivation is also widely practiced by the farmers.

The importance of vesicular-arbuscular (VA) mycorrhiza in agricultural crops is widely recognised (Mosse, 1973; Gerdemann, 1975; Mishra and Verma, 1982). Paddy has been reported to be infected by VA mycorrhizal fungi under field conditions (Iqbal et al., 1975; Manjunath et al., 1981b). Bhattarai (1983) observed high VA mycorrhizal infection in different paddy varieties under upland cultivation practice. Growth response of paddy plants to VA mycorrhizal infection was found by Iqbal et al., (1978a). Still, however, detailed study on the role of VA mycorrhiza in paddy is awaited.

Phosphorus is by far considered as the most important nutrient concerned in growth responses of plants to VA mycorrhiza (Mosse, 1973; Krishna and Bagyaraj, 1982). The improved phosphorus nutrition of VA mycorrhizal plants results from an increased efficiency in phosphorus uptake from the soil (Sanders and Tinker, 1971; Powell, 1975a; Gianinazzi Pearson et al., 1981). There is now substantial evidence (Rhodes and Gerdemann, 1975; Owusu Bennoah and Wild, 1979; Graham et al., 1982b) to show that increased phosphorus uptake is possible because of the capacity of the external

hyphae of VA mycorrhizal fungi to absorb phosphate from the soil and translocate it into the host root. This enables the plants to have a longer, better distributed surface for absorption. Besides P, the VA mycorrhizal fungi also enhance uptake of micro-elements (Rhodes and Gerdemann, 1978; Zaag et al., 1979; Bagyaraj and Manjunath, 1980; Jensen, 1982).

Different strains of VA mycorrhizal fungi vary in their efficiency for nutrient uptake (Jensen, 1982; Davis and Young, 1985). This difference is thought to be due to the difference in ability of extension of hyphal distances travelled from the root surface (Abbott and Robson, 1985b). The cultivars of the same species have also been reported to vary in their ability to form mycorrhiza (Azcon and Ocampo, 1981; Manjunath et al., 1981b; Gueye, 1983).

VA mycorrhizal infection and endogonaceous spore population were found to be reduced by addition of soluble phosphate (Abbott and Robson, 1977; Asimi et al., 1980; Jensen and Jakobsen, 1980). Menge et al., (1978) proved that concentration of phosphorus within the plant root and not the soil phosphate status was the regulatory factor in determining colonization, infection and spore production by VA mycorrhizal fungi. Addition of nitrogen fertilizer in soil was also reported to decrease spore production and VA mycorrhizal infection (Hayman, 1970; Chambers et al., 1980), though

application of increasing amounts of nitrogen in soil was also found to increase VA mycorrhizal infection (Hepper, 1983).

Moisture level of soil and frequency of irrigation affect VA mycorrhizal status in host root (Levy et al., 1983; Anderson et al., 1986). VA mycorrhizal fungi were reported to impart resistance to the host against drought stress (Levy and Krikun, 1980; Hardie and Leyton, 1981), though there are some reports which do not agree with this hypothesis (Hetrick et al., 1984). Most reports now accept that the growth response of plants to VA mycorrhiza under drought stress is due to improved P nutrition and more extensive soil water extraction (Allen, 1982; Busse and Ellis, 1985).

Rhizospheric micro-organisms were found to interact with VA mycorrhizal fungi. Bagyaraj and Menge (1978) observed an increased population of bacteria and actinomycetes in the rhizosphere of tomato inoculated with Glomus fasciculatum. A beneficial effect on plant growth was achieved by Manjunath et al., (1981a) when Glomus fasciculatum was inoculated in combination with Aspergillus niger. Verma (1982) also reported increased growth in Eupatorium adenophorum when Glomus tenue was inoculated in combination with Trichoderma viridae. T. viridae also increased mycorrhizal infection. As far as soil borne fungal pathogens are concerned, variable reports are available. Davis et al., (1978) observed more

susceptibility in mycorrhizal roots of avocado against Phytophthora cinnamomi but Matare and Hattingh (1978) could not find any effect of Glomus fasciculatum on avocado root rot caused by P. cinnamomi. Many reports are available which advocate that VA mycorrhizal fungi confer resistance to host against root pathogens (Schonbeck, 1978; Dehne and Schonbeck, 1979 a; Chakravarty and Mishra, 1986 a & b). Different VA mycorrhizal fungi were also reported to confer variable tolerance or resistance to the host against pathogen (Davis and Menge, 1981).

In Meghalaya, where most of the farmers do not use the fertilizers due to lack of efficient transportation facilities and also due to their socio-economic conditions, VA mycorrhizal fungi can play an important role in nutrient uptake. The local climate is also very conducive for disease development. Keeping these points in view, a study on the role of VA mycorrhizal fungi on paddy was planned as enumerated below :-

1. Status of VA mycorrhizal fungi of paddy in different varieties and under different cultivation practices.
2. Effect of VA mycorrhiza on growth and phosphorus uptake in different paddy varieties under different soil phosphorus levels.
3. Effect of VA mycorrhiza on growth and nutrient uptake in paddy under different soil nitrogen and phosphorus levels.

4. Effect of VA mycorrhiza on growth of paddy under different water stress conditions.
5. Interaction of VA mycorrhizal fungi with rhizospheric fungi of paddy.

REVIEW OF LITERATURE